

Evaluation of Application Methods Efficiency of Zinc and Iron for Canola (*Brassica napus* L.)

Ahmad BYBORDI, Garib MAMEDOV

Baku State University, Faculty of Biology, 23 Z. Khalilov Street, 370148 Baku, Azerbaijan; abaybordy@yahoo.com

Abstract

In order to evaluation of application method efficiency of zinc and iron microelements in canola, an experiment was conducted in the Agricultural Research Station of Eastern Azerbaijan province in 2008. The experimental design was a RCBD with eight treatments (F₁: control, F₂: iron, F₃: zinc, F₄: iron + zinc in the form of soil utility, F₅: iron, F₆: zinc, F₇: iron+ zinc in the form of solution foliar application, and F₈: iron + zinc in the form of soil utility and foliar application). Analysis of variance showed that there were significant differences among treatments on given traits, antioxidant enzymes activity, fatty acids percentage, plant height, seed weight to capitulum weight ratio, protein percentage, oil percentage, oil yield, 1000 seed weight, seed yield, nitrogen, phosphorous and potassium percentage of leaves, zinc and iron content of leaves and capitulum diameters. The highest seed yield, oil yield, oil percentage, 1000 seed weight, seed weight to capitulum weight ratio and protein percentage were obtained from the soil and foliar application of iron + zinc treatments (F₈). Also, the highest amounts of nitrogen, phosphorous and potassium concentration in leaves were achieved from control treatment which was an indication of non-efficiency of iron and zinc on the absorption rate of these substances in the leaves. The correlation between effective traits on the seed yield, such as, capitulum diameter, number of seed rows in capitulum, seed weight to capitulum weight ratio and 1000 seed weight were positively significant. In general, foliar and soil application of zinc and iron had the highest efficiency in aspect of seed production. The comparison of the various methods of fertilization showed that foliar application was more effective than soil application. Also, micronutrient foliar application increased concentration of elements, especially zinc and iron. Antioxidant enzymes activity was different in response to treatments also the highest palmitoleic, oleic and myristic acid were observed in F₆ and F₇ treatments.

Keywords: foliar application, iron, soil utility, canola, zinc

Introduction

Canola (*Brassica napus* L.) is grown in different agro-climatic zones of the world, differing in soil nutrient status. The use of foliar fertilizing in agriculture has been a popular practice with farmers since the 1950s, when it was learned that foliar fertilization was effective and economic. Recent research has shown that a small amount of nutrients, particularly Zn, Fe and Mn applied by foliar spraying increases significantly the yield of crops (Sarkar *et al.*, 2007; Wissuwa *et al.*, 2008). Also, foliar nutrition is an option when nutrient deficiencies cannot be corrected by applications of nutrients to the soil (Crabtree, 1999; Sarkar *et al.*, 2007; Cakmak, 2008). It is likely therefore, in open-field conditions, where the factors that influence the uptake of the nutrients are very changeable, foliar fertilization can get considerable importance. Among the micro-nutrients, Zn and Fe nutrition can affect the susceptibility of plants to drought stress (Sultana *et al.*, 2001; Khan *et al.*, 2003; Cakmak, 2008).

Zn is known to have an important role either as a metal component of enzymes or as a functional, structural or reg-

ulatory cofactor of a large number of enzymes (Grotz and Guerinot, 2006). Zinc also plays an important role in the production of biomass (Kaya and Higgs, 2002; Cakmak, 2008). Furthermore, zinc may be required for chlorophyll production, pollen function and fertilisation (Kaya and Higgs, 2002; Pandey *et al.*, 2006). Low solubility of Zn in soils rather than low total amount of Zn is the major reason for the widespread occurrence of Zn deficiency problem in crop plants (Cakmak, 2008). High seed-Zn has very important physiological roles during seed germination and early seedling growth (Cakmak, 2008). The review by Cakmak (2008) provides further reasons and relevant research for benefits of high seed-Zn on plant growth. As an activator of CuZn- or Mn-SOD, zinc or manganese is involved in membrane protection against oxidative damage through the detoxification of reactive oxygen species (Marschner, 1995).

Iron plays essential roles in the metabolism of chlorophylls. External application of Fe increased photosynthesis, net assimilation and relative growth in seawater-stressed rice (Sultana *et al.*, 2001). This is especially true for soils of high pH where equilibrium conditions favour the oxida-

tion of plant-available Fe^{+2} to unavailable Fe^{+3} . Plant yield on many soils is, therefore, limited by poor Fe availability, rather than a low Fe content in the soil. Also Fe leaching is the main pathway for Fe loss in coarse-textured soil with high pH, while excessive Fe uptake was the main pathway for Fe loss in clay-textured and acid soil. Application of Zn or Fe has been reported significant positive effects, in most cases, on growth measurements and chemical composition of safflower (Lewis and McFarlane, 1986), lupine (Brennan *et al.*, 2001), cumin (El-Sawi and Mohamed, 2002), soybean (Gadallah, 2000; Heitholt *et al.*, 2002), barley (Genc *et al.*, 2004; Hebborn *et al.*, 2005), wheat (Lu *et al.*, 2004), sunflower (Mirzapour and Khoshgoftar, 2006), mustard (Chatterjee and Khurana, 2007), common bean (Fernandes *et al.*, 2007) and rice (Wissuwa *et al.*, 2008). Even though iron is one of the most abundant elements in soils, the low solubility product of iron minerals makes the inorganic form of iron unavailable to plants and forms the most common widespread nutritional disorder world over (Welch *et al.*, 1991). Plants subjected to iron deficiency excess respond in different ways (Abadia, 1992). It is believed that under conditions of iron stress, some plants can increase their absorption capacity for iron. The mechanism affecting the acquisition of this essential microelement is often present in aerobic soils in the form of Fe (Marschner, 1995).

However, to the best of our knowledge, information regarding application method efficiency of zinc and iron on the growth and development of canola is not available. Therefore, the purpose of this study is to understand whether application of micronutrients as foliar application is better or soil application.

Materials and methods

The experiment was conducted on the research farm of Agricultural Research Station of Eastern Azerbaijan province, Azerbaijan, Iran ($37^{\circ}58'$ N latitude, $46^{\circ}3'$ E longitude and altitudes of 1320 m) during 2008 growing seasons. Site of study has cold winter and warm summer. The yearly average precipitation (30-years long term period) which is mostly occurred during the spring months is 273.1 mm. The mean annual temperature was 11°C .

Soil sampling and analysis

Prior to the beginning of experiment, soil samples were taken in order to determine the physical and chemical properties. A composite soil samples were collected at a depth of 0-30 cm. It was air dried, crushed, and tested for

physical and chemical properties. The research field had a loamy soil. Details of soil properties are shown in Tab. 1.

Field preparation and applying the treatments

After plough in fall and two disks in spring, the land was flatted by leveller and then plots were prepared. The experimental design was laid out in a Randomized Complete Block Design with four replications. The plots had 4 m length and 3 m width consisted of five rows, 0.6 m apart. Between all plots, 1 m distance was kept to eliminate all influence of lateral water movement. According to results of soil analysis 50 k.ha^{-1} super phosphates triple and 300 kg.ha^{-1} urea was used. All of super phosphates triple and one third of urea were distributed in plots and mixed with surface soil before seed sowing. Rest of urea was used when plants have 6 or 8 leaves and at flowering stage as topdress. Different treatments of micronutrient soil application which were including: 25 k.ha^{-1} iron from secostriene and 40 kg.ha^{-1} from zinc sulphate were distributed in plots and mixed with surface soil before seed sowing.

Seed sowing and irrigation

The canola seeds (*Brassica napus* L. C.V. 'SLM₀₄₆') were sown on 15 July. The between row distance and within distance were 60 cm and 25 cm, respectively. The flooding Irrigation was carried out as similar in all of plots during study. Two week after sowing weed control was efficiently performed by hand. Micronutrient foliar application was done at two times, one time when plants had 6 or 8 leaves and another when they have 10 or 12 leaves (early of flowering stage). The iron and zinc were sprayed on plants with concentrate of 2 parts per thousand and 3 part per thousand, respectively. In control treatments plants were sprayed by water. Description of all treatments is shown in Tab. 2.

Plant sampling, harvesting and scrutiny

At early flowering, number of leaves and leaf area were counted. Leaf area was estimated by multiplying of length and width of leaves in 0.68 coefficients (Rao and Saran, 1991). Also, fresh leaves samples were taken of similar leaves. The samples were frozen in liquid nitrogen and stored at -80°C until biochemical analysis.

After that, at flowering stage four upper leaves were taken for determine of N, P, K, Mn, Fe and Zn content of leaves. The samples were washed by distilled water and dried in oven at 70°C during 48 h. Total N, P, K, Mn, Fe and Zn content were determined through atomic absorp-

Tab. 1. Physical and chemical soil properties

	Sand	Silt	Clay	Texture	Calcite	O.M	EC
Depth	42%	28%	30%	Loamy	18%	0.32%	2.6 d.m^{-1}
0-30 cm	N	P	K	Mn	Fe	Zn	Cu
	0.03%	14 mg.Kg^{-1}	320 mg.Kg^{-1}	2.8 mg.Kg^{-1}	2.2 mg.Kg^{-1}	0.22 mg.Kg^{-1}	1.8 mg.Kg^{-1}

Tab. 2. Different treatments of micronutrient application

F ₁	Control	Water
F ₂	Iron (soil application)	25 k.ha ⁻¹ iron from secostrine
F ₃	Zinc (soil application)	40 k.ha ⁻¹ zinc sulphate
F ₄	Iron + Zinc (soil application)	25 k.ha ⁻¹ iron from secostrine + 40 k.ha ⁻¹ zinc sulphate
F ₅	Iron (foliar application)	2 parts per thousand
F ₆	Zinc (foliar application)	3 parts per thousand
F ₇	Iron + Zinc (foliar application)	2 parts per thousand + 3 parts per thousand
F ₈	Iron + Zinc (soil and foliar application)	25 k.ha ⁻¹ iron from secostrine + 40 k.ha ⁻¹ zinc sulphate + 2 parts per thousand + 3 parts per thousand

tion method (Perkinelmer 1012, USA). Plants height was measured at end of flowering stage by five sample plant.

At the physiological maturity stage plants were harvested and capitulum diameter, row number in capitulum, number of seed in capitulum, seed weight to capitulum weight ratio, 1000 seed weight, seed yield, total oil percentage and total protein percentage and oil yield were measured.

The fatty acid compositions of the canola seed oils were determined by gas chromatography (GC) (Metcalf *et al.*, 1966). The contents of palmitoleic, linolenic, oleic and myristic acids were determined using a computing integrator. The effects of the independent variables on oil content and palmitoleic, Linolenic, oleic and myristic acid concentrations of the oil were analyzed on a percentage basis.

Oil percentage and protein percentage were measured by soxhlet and Kjelttec method, respectively. Oil yield was calculated via product seed yield in percentage oil.

Antioxidant enzyme activity

Peroxidase

Peroxidase activity was determined by the oxidation of guaiacol in the presence of H₂O₂. The increase in absorbance was recorded at 470 nm (Ghanati *et al.* 2002). The reaction mixture contained 100 µL crude enzyme, 500 µL H₂O₂ 5 mm, 500 µL guaiacol 28 mm and 1900 µL phosphate buffer 60 mm (pH 7.0).

Catalase

Catalase activity was estimated by the method of Cakmak and Horst (1991). The reaction mixture contained 100 µL crude enzyme extract, 500 µL 10 mm H₂O₂ and 1400 µL 25 mm phosphate buffer. The decrease in the absorbance at 240 nm was recorded for 1 min.

Glutathione reductase

Glutathione reductase activity was determined by the method of Halliwell and Foyer (Halliwell and Foyer, 1978). Its activity was assayed in a 1 mL reaction mixture containing 0.25 mL of 100 mM potassium phosphate buffer (pH 7.0), 0.05 mL of 10 mM oxidized glutathione, 0.12 mL of 1mM NADPH, 0.48 mL of distilled water, and 0.1 mL of enzyme extract. The resultant decrease in NADPH was observed at 340 nm.

Superoxide dismutase

The activity of SOD was measured according to the method of Giannopolities and Ries (1977). The assay medium contained 50 mM phosphate buffer (pH 7.8), 13 mM methionine, 75 mM p-nitro blue tetrazolium chloride, 2 mM riboflavin, 0.1 mM EDTA and 5 mL enzyme extract. Glass test tubes containing the assay medium were illuminated with a fluorescent lamp (120 W); identical tubes that were not illuminated served as blanks. After illumination for 15 min, the absorbance was measured at 560 nm. One unit of enzyme activity was determined as the amount of the enzyme to reach an inhibition of 50% nitro blue tetrazolium reduction rate by monitoring the absorbance at 560 nm.

Reduced glutathione

The reduced glutathione content was assayed as described by Griffith and Meister (1979). 200 mg of fresh material was ground with 2ml of 2% metaphosphoric acid and centrifuged at 17,000 rpm for 10 min. Adding 0.6 ml 10% sodium citrate neutralized the supernatant. One milliliter of assay mixture was prepared by adding 100 µl extract, 100 µl distilled water, 100 µl of 6mM 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB) and 700 µl of 0.3mM NADPH. The mixture was stabilized at 25 °C for 3–4 min. Then 10 µl of glutathione reductase was added and read the absorbance at 412 nm in spectrophotometer.

Statistical analysis of data

All data were analyzed from analysis of variance (ANOVA) using the MSTAT-C. Least significant difference test was used to measure statistical differences between treatment methods and controls. Also the charts were drawn by Excel, Microsoft Office 2003.

Results and discussion

The results of analysis of variance showed that, effect of micronutrients was significant on all traits except number of leaves in plant, leaf area, number of seed row in capitulum, seed number in capitulum, kernel to seed percentage and Mn content (Tab. 3 and 4).

Tab. 3. Effect of micronutrient application on some agronomical traits of canola

S.O.V	d.f	PH	LN	LA	CD	SW	PP	OP
Block	3	110.78	18.58	0.53	1.40	65.56	0.04	2.54
Treatment	7	97.19**	7.30	0.11	2.17*	59.19**	3.37**	13.69**
error	21	20.53	3.88	0.15	1.06	6.33	0.005	0.49
C.V		3.51	9.20	23.54	5.45	3.71	0.34	1.62
S.O.V	d.f	KS	OY	SY	LN	LZ	LM	
Block	3	3.03	12664.27	42443.65	0.11	28.53	261.69	
Treatment	7	3.89	155428.34**	489700.60**	0.16**	256.56**	9.81	
error	21	4.59	9975.70	53768.74	0.01	16.22	14.69	
C.V		2.80	4.60	4.64	4.35	14.30	5.45	

*, ** significant at the 0.05 and 0.01 probability levels, respectively

PH: plant height; LN: leaf number; LA: leaf area; CD: capitulum diameter; SW: seed weight; PP: protein percentage; OP: oil percentage; Kernel to seed percentage; OY: oil yield; SY: seed yield; LN: leaf nitrogen; LP: leaf phosphorous; LK: leaf potassium; LI: leaf iron; LZ: leaf zinc; LM: leaf manganese

Antioxidant enzyme activity

The highest POD activity was observed in F₆ and F₈ treatment while the lowest activity was related to F₂ treatment (Tab. 5). There was not significant difference between F₁ and F₃ and between F₅ and F₇ treatments (Tab. 5).

It seems that Zn foliar application causes increase of POD activity because soil application had not additive effect on activity of this enzyme in contrast Fe application decreased POD activity. Our findings were in agreement with the results reported by Jiang and Huang (2001) and Habibi *et al.* (2004). The simultaneous increase in the activity of these enzymes contributes to a decrease of the deleterious effects of H₂O₂ under stress.

Control treatment had the highest CAT activity (Tab. 5). The lowest CAT activity was achieved from F₄, F₆ and F₈ treatments, as among these treatments were not significant difference (Tab. 5). Similarly, there was no difference between F₃ and F₅ application method. Fe and Zn application decreased CAT activity, due to scavenging of reactive oxygen species by other antioxidant enzymes. These results were agreement with the results reported by Bailly *et al.* (2000).

Glutathione reductase was affected by fertilizers too. As F₂ and F₃ treatments had the highest enzyme activity while the lowest was observed from F₈ treatment (Tab. 5). There was not significant difference between F₄ and F₆ treatments too (Tab. 5). The highest and the lowest GSH

activity were related to F₁, F₂, F₃ and F₆, F₇, F₈ treatments, respectively (Tab. 5).

The results showed that, F₇ and F₈ treatments increased significantly SOD activity. The lowest activity was observed in F₁, F₃ and F₅ treatments (Tab. 5). Cakmak (2000) reported that Zn deficiency may inhibit the activities of a number of antioxidant enzyme.

It has been demonstrated that environmental stress induces oxidative stress in plant tissues. Exposes chloroplasts to excessive excitation energy, may increase generation of reactive oxygen species and induce oxidative stress. To overcome the effects of oxidative stress, plants make use of a complex antioxidant system. Relatively higher activities of reactive oxygen species scavenger enzymes have been reported in many stressed plants, which suggest that the antioxidant system plays an important role in plants against environmental stresses (Habibi *et al.*, 2004).

Superoxide dismutase may function as a reactive oxygen species scavenger, by converting O₂⁻ to H₂O₂ (Baily *et al.* 2000). Hydrogen peroxide is converted to oxygen and water by CAT and POX. Even though high SOD activity protects plants against superoxide radicals, it can not be considered solely responsible for membrane protection against peroxidation. In general, micronutrient application had different effects on antioxidant enzymes activity, in some cases increase and in some cases decrease was observed.

Tab. 4. Effect of micronutrient application on antioxidant enzyme activity and fatty acid percentage of canola

S.O.V	df	POD	CAT	GR	GSH	SOD	Palmitoleic acid	Linolenic acid	Oleic acid	Myristic acid
Block	2	2.16	2.16	0.00	0.00	3.51	1.29	0.64	0.37	0.16
Treatment	7	30.76**	13.94**	0.01**	0.01**	24.49**	22.47**	9.84**	18.19**	3.47**
Error	14	0.97	0.50	0.00	0.00	0.24	4.38	1.34	1.42	0.43
C.V		3.10	3.58	2.63	3.46	3.53	9.89	17.36	1.76	16.66

*, ** significant at the 0.05 and 0.01 probability levels, respectively

POD: peroxidase, CAT: Catalase; GR: glutathione reductase; SOD: superoxide dismutase

Tab. 5. Comparison of means on antioxidant enzyme activity and fatty acid percentage of canola

Treatments	POD	CAT	GR	GSH	SOD	palmitoleic acid	Linolenic acid	Oleic acid	Myristic acid
F ₁ control	30.66cd	24.00a	0.46b	0.73a	10.50d	17.66c	3.7e	61.66b	2.60c
F ₂ Iron (soil application)	26.33e	21.33b	0.52a	0.75a	12.33c	20.00bc	6.00cd	67.33a	2.93bc
F ₃ Zinc (soil application)	29.00d	20.00c	0.51a	0.71a	11.00d	22.33ab	8.00abc	68.66a	3.40bc
F ₄ Iron + Zinc (soil application)	31.00c	18.00d	0.40d	0.65b	13.00c	17.00c	8.66ab	68.66a	3.66bc
F ₅ Iron (foliar application)	33.00b	19.33c	0.43c	0.64b	15.33b	22.00ab	9.00a	68.00a	4.20b
F ₆ Zinc (foliar application)	36.00a	17.66d	0.40d	0.59c	15.00b	25.00a	5.00de	69.66a	5.50a
F ₇ Iron + Zinc (foliar application)	34.00b	19.66c	0.36e	0.58c	18.00a	22.33ab	6.66bcd	68.33a	5.50a
F ₈ Iron + Zinc (soil and foliar application)	34.66ab	17.66d	0.32f	0.62bc	17.66a	23.00ab	6.33cd	67.66a	4.03b

For a given means within each column followed by the same letter are not significantly differences ($p < 0.05$)




Fig. 1. Changes of protein percentage due to different methods of iron and zinc application. F₁ = control, F₂ = Iron (soil application), F₃ = Zinc (soil application), F₄ = Iron + Zinc (soil application), F₅ = Iron (foliar application), F₆ = Zinc (foliar application), F₇ = Iron + Zinc (foliar application), F₈ = Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences ($p < 0.05$)




Fig. 2. Changes of oil percentage due to different methods of iron and zinc application. F₁ = control, F₂ = Iron (soil application), F₃ = Zinc (soil application), F₄ = Iron + Zinc (soil application), F₅ = Iron (foliar application), F₆ = Zinc (foliar application), F₇ = Iron + Zinc (foliar application), F₈ = Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences ($p < 0.05$)




Fig. 3. Changes of 1000 seed weight due to different methods of iron and zinc application. F₁ = control, F₂ = Iron (soil application), F₃ = Zinc (soil application), F₄ = Iron + Zinc (soil application), F₅ = Iron (foliar application), F₆ = Zinc (foliar application), F₇ = Iron + Zinc (foliar application), F₈ = Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences ($p < 0.05$)

Fatty acids

Analysis of fatty acids by GC showed that, the application of Zn and Fe increased the fatty acids compared to




Fig. 4. Changes of plant height due to different methods of iron and zinc application. F₁ = control, F₂ = Iron (soil application), F₃ = Zinc (soil application), F₄ = Iron + Zinc (soil application), F₅ = Iron (foliar application), F₆ = Zinc (foliar application), F₇ = Iron + Zinc (foliar application), F₈ = Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences ($p < 0.05$)

the control (Tab. 5). In case of palmitic, palmitoleic and myristic acid foliar application was better than soil application while regarding linolenic acid soil application had better effect. Between foliar application and soil application was not difference on oleic acid content but applica-




Fig. 5. Changes of leaf iron due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)




Fig. 6. Changes of leaf potassium due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)




Fig. 7. Changes of leaf nitrogen due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)




Fig. 8. Changes of leaf phosphorous due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)




Fig. 9. Changes of seed yield due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)




Fig. 10. Changes of oil yield due to different methods of iron and zinc application. F₁= control, F₂= Iron (soil application), F₃= Zinc (soil application), F₄= Iron + Zinc (soil application), F₅= Iron (foliar application), F₆= Zinc (foliar application), F₇= Iron + Zinc (foliar application), F₈= Iron + Zinc (soil and foliar application). For a given means within each column followed by the same letter are not significantly differences (p < 0.05)

tion of these elements increased significantly oleic content than control treatment (Tab. 5). Zakaria *et al.* (2001) concluded that Zn foliar application significantly increased the total unsaturated fatty acids in cotton.

The highest palmitoleic acid was related to F₃, F₅, F₆, F₇ and F₈ treatments also, the highest and the lowest linolenic acid percentage were achieved from F₃, F₄, F₅ and control treatment, respectively (Tab. 5).

The highest myristic percentage was achieved from F₆ and F₇ treatments (Tab. 5). It was observed that control treatment had the lowest myristic percentage (Tab. 5).

Canola oil consists of different types of saturated and unsaturated fatty acids (palmitic acid, stearic acid, oleic acid, linoleic acid etc.). The palmitic acid and stearic acid are the major saturated fatty acids, whereas oleic and linoleic acids are unsaturated. Fatty acid composition of canola in particular and other oil seed crops in general, are influenced by fertilizing management. The findings of present study are shows that fatty acid composition is affected by iron and zinc application, for example, Kheir *et al.* (1991) found that the higher N-rate increased the percentage of unsaturated fatty acids and decreased saturated fatty acids of flax oil.

Protein percentage

The highest and the lowest protein percentage were observed in F₈ treatment and control, respectively. Kaya and Higgs (2002) pointed that the application of Zn fertilizers significantly increased the protein content of tomato. But it was in consistent with results of Singh, S. (2000) which suggest that protein content does not affected by fertilizers treatment and is related to genotypes. Cakmak *et al.* (1989) stated that under condition of zinc deficiency protein content decreases. Iron and zinc are two important elements in enzymes structure involved in amino acid biosynthesis and thus amino acids are the base of protein synthesis, protein content increases in use of these micronutrients. Our results are agreed with results of Zakaria *et al.* (2001) on cotton.

Oil yield and percentage

The F₂, F₄ and F₈ treatments produced the highest oil percentage. The lowest oil percentage was obtained from foliar application treatment and control treatment. It seems that, soil application of micronutrients is more benefit to oil biosynthesis. Singh and Sinha (2005) reported the decrease in oil concentration may be due to oxidation of some polyunsaturated fatty acids. The highest and the lowest oil yield were achieved from F₈ and control treatments. In general, there was not significant different between micronutrient application method. Because of direct relation between seed yield and oil yield, increase in seed yield causes in higher oil yield (Singh, S. 2000).

1000 seed weight

The highest 1000 seed weight was observed in F₈ treatment and other foliar application treatments. Zinc is required for the biosynthesis of the plant growth regulator such as IAA and for carbohydrate and N metabolism which leads to high yield and yield components. This may be due to provision of macro and micro nutrients at latter stages which might have enhanced accumulation of assimilate in seeds and thus resulting in heavier seeds.

Seed yield

The highest seed yield was produced from F₈ treatment. Masoni *et al.* (1996) showed that, zinc increases seed yield. Heitholt *et al.* (2002) reported that seed yield of soybean increased while Cu, Mn, Zn, and Fe applied individually. The increase in yield with soil applied Fe might have been the result of increase in the amount of available Fe in soil. The increase in seed yield with foliar application of Fe could be attributed to the direct absorption of the element by the foliar sprayed with Fe.

In proceed to the previous studies (Cakmak, 1999), all methods of Zn application for plants significantly increased grain yield. Micronutrients increases photosynthesis rate and improves leaf area duration thus seed yield will be increased. Zinc plays important role in tryptophan biosynthesis, later is precursor of auxine also zinc is founded in phosphoenolpyruvate carboxylase structure. Another element that is iron is necessary to chlorophyll synthesis and its critical element in electron transport chain in photosystems. Iron deficiency leads to many disorders in chloroplasts. Ferredoxin is an important iron-containing protein involved in electron-transfer. Singh *et al.* (1975) observed that, zinc increases canola significantly seed yield. The lowest seed yield was obtained from control treatment.

Leaf nitrogen content

The results showed that, micronutrient application decreased leaf nitrogen content. The highest nitrogen content was related to control treatment while the lowest content was obtained from F₈ treatment. Foliar application had more effect on nitrogen content reduction. It seems that, decrease in nitrogen content is due to competitive effect between elements such as zinc and nitrogen.

Leaf phosphorous content

Similar upon results, micronutrient fertilizing decreased phosphorous content in leaves. These results were inconsistent with that obtained from Movchan and Sobornikova (1972). The highest phosphorous content was related to control treatment. Zinc decreases phosphorous content because there is competitive effect between phosphorous and zinc in uptake of ions. Sahota and Arora (1981) reported that there was no significant effect of Zn on P content of leaves.

Leaf potassium content

There was significant decrease in potassium content of plants which were treated with iron or zinc than control plants. The lowest potassium content was observed in F₈ treatment. It shows that, iron and zinc decreases potassium content in leaves of canola. Stoyanova and Doncheva (2002) remarked that the K content of the stems and leaves was not significantly affected by different concentrations of Zn.

Leaf iron content

The highest iron contents were observed in four treatments, F₈, F₆, F₅ and F₄, respectively. The application of Fe through soil as well as foliar application caused a marked increase in the total content of Fe in the fenugreek plants. Whereas in foliar application method iron absorption is faster and easier than soil application the highest iron content was observed in these treatments. In soil application due to organic matters banding with chemical fertilizers that have high ability to absorb and hold nutrients, and positioning these substances near hairy roots, results in better availability to plant and thus causing in higher iron content. Studying previous researches show that best results achieved in using iron sulphate (Kalbasi *et al.* 1998). And may be the use of zinc sulphate causes better absorption in iron. Iron deficiency leads to chlorophyll degradation and chlorosis. It's reported that, iron is an essential element in protein synthesis and iron deficit decreases plant growth (Agarwala *et al.*, 1965). Also, iron is involved electron transport in photosystems. Already decrease of chlorophyll content due to iron deficit was reported by Masoni *et al.* (1996).

Leaf zinc content

The highest zinc content was observed in F₆, F₇ and F₈ treatments, respectively. Foliar application of zinc increased zinc content more than soil application. Zinc uptake is easier from leaves in compare soil application. Zinc uptake is lower in lime soils. Grawel and Graham (1999) have been reported that, zinc application increases zinc concentration in seed, root and leaves. Increase in Zn concentration reported in soil application of S and Fe. These results were in agreement with those reported by El-Gazzar *et al.* (1979) and Foregoni *et al.* (1974). Zinc is essential element in enzymatic system such as superoxide dismutase enzyme. Zinc plays important role in auxine and protein synthesis and it is essential for seed setting.

Conclusions

Iron and zinc uptake are controlled by the two major factors, availability of these elements in the soil and the ability of plants to acquire them. Application methods of micronutrients are very important to attain the best ab-

sorption. Sometimes response of the plants is different to application methods of fertilizers, for example in calcareous soil Fe and Zn are not available for plants, in this times, foliar application is a useful method for nourish of the plants. The results of this study demonstrated that, Fe and Zn had positive effect on yield and quality of canola oil. In addition to this, we also suggest that Ca and Fe should be sprayed on plants to reach the best quality and quantity in canola production.

References

- Abadia, J. (1992). Leaf responses to Fe deficiency: a review. *J. Plant Nutr.* 15:1699-1713.
- Agarwala, S. C., C. P. Sharma and S. Farooq (1965). Effect of iron supply on growth, chlorophyll, tissue iron and activity of certain enzymes in maize and radish. *Plant Physiol.* 40:493-499.
- Balakrishnan, K. (2000). Peroxidase activity as an indicator of the iron deficiency banana. *Ind. J. Plant Physiol.* 5:389-391.
- Bailly, C., A. Benamar, F. Corbineau and D. Come (2000). Antioxidant systems in canola seeds as affected by priming. *Seed Science Research* 10, 2: 35-42. Bailly, C., Benamar, A., Corbineau, F. and D. Come (2000). Antioxidant systems in canola seeds as affected by priming. *Seed Science Research* 10, 2: 35-42.
- Brennan, R. F., M. D. A. Bolland and G. Shea (2001). Comparing how *Lupinus angustifolius* and *Lupinus luteus* use zinc fertilizer for seed production. *Nutr. Cycl. Agroecosyst.* 59: 209-217.
- Cakmak, I., H. Marschner and F. Bangert (1989). Effect of zinc nutritional status on growth, protein metabolism and levels of indole-3-acetic acid and other phytohormones in bean (*Phaseolus vulgaris* L.). *J. Exp. Bot.* 40: 404-412.
- Cakmak, I. (2008). Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant Soil.* 302: 1-17.
- Cakmak, I. and W. Horst (1991). Effect of aluminium on lipid peroxidation, superoxide dismutase, catalase and peroxidase activities in root tip of soybean (*Glycine max*). *J. Plant Physiol.* 83:463-468.
- Cakmak, I., M. Kalayci, H. Ekis, J. Brauni, Y. Kilinc and A. Yilmaz (1999). Zn deficiency as a practical problem in Plant and human nutrition in Turkey: a NATO-science for stability project. *Field Crop Res.* 60: 175-188.
- Chatterjee, C. and N. Khurana (2007). Zinc stress-induced changes in biochemical parameters and oil content of mustard. *Commun. Soil Sci. Plant Anal.* 38:751-761.
- Crabtree, W. L. (1999). Deep placement of Mn fertiliser on a sandy soil increased grain yield and reduced split seed in *Lupinus angustifolius*. *Plant Soil.* 214:9-14.
- Deshmukh, M. R. and K. N. Wavhal (1999). Effects of iron on growth and flowering of aster. *J. Maha Agric. Univ.* 23: 99-101.

- El-Gazzar, A. M., A. Wallace and A. M. Rokba (1997). Growth and Leaf Mineral Composition of Oranges, Olive, Plums and Grapes as Influenced by Calcium Carbonate Addition to the Soil in a Greenhouse. *Egypt J. Hort. Sci.* 2:141-149.
- El-Sawi, S. A. and M. A. Mohamed (2002). Cumin herb as a new source of essential oils and its response to foliar spray with some micro-elements. *Food Chem.* 77:75-80.
- Eppendorfer, W. H. and B. O. Eggum (1994). Dietary fibre, starch, aminoacids and nutritional value of potatoes as affected by sulfur, nitrogen, phosphorus, calcium and water stress. *Acta Agric. Scand. Sect. B, Soil Plant Sci.* 44:107-115.
- Fernandes, D. S., R. P. Soratto, S. M. Kulczynski, G. A. Biscaro and C. J. Dos Reis (2007). Yield and physiological quality of common bean seeds as affected by manganese foliar application. *Pesq. agropec. bras. Brasília* 42:419-426.
- Foregoni, M., M. Boselli, C. Bartoletti and G. Dorotea (1984). Effect of Iron Deficiency or an Excess of Leaf-applied Iron on the Mineral and Biochemical Composition and on the Production Parameters of Chlorotic Grapevines. *Connassiance de la vinge et due Vin, Hort. Abst.* 54:95-110.
- Gadallah, M. A. A. (2000). Effects of indole-3-acetic acid and zinc on the growth, osmotic potential and soluble carbon and nitrogen components of soybean plants growing under water deficit. *J. Arid Environ.* 44: 451-467.
- Genc, Y., G. K. McDonald and R. D. Graham (2004). Differential expression of zinc efficiency during the growing season of barley. *Plant Soil* 263:273-282.
- Ghanati, F., A. Morita and H. Yokota (2002). Induction of suberin and increase of lignin content by excess Boron in Tobacco cell. *Soil Sci. Plant Nutr.* 48, 3:357-364
- Giannopolities, C. and S. Ries (1977). Superoxide dismutase. Part I: occurrence in higher plants. *J. Plant Physiol.* 59:309-314.
- Griffith, O. W. (1979) Potent and specific inhibition of glutathione synthesis by buthionine sulfoximine (s-n-butylhomocysteine sulfoximine). *J. Biol. Chem.*, 254: 7558-7560.
- Grawel, H. S. and R. Graham (1999). Residual effect of subsoil zinc and oilseed rape genotype on the grain yield and distribution of zinc in wheat. *Plant and Soil* . 207:29-36.
- Goos, R. J., B. Johnson, G. Jackson and G. Hargrove (2004). Greenhouse evaluation of controlled release iron fertilizers for soybean. *J. Plant Nutr*:27, 43
- Grotz, N. and M. L. Guerinot (2006). Molecular aspects of Cu, Fe and Zn homeostasis in plants. *Biochim. Biophys. Acta* 1763: 595-608.
- Heitholt, J. J., J. J. Sloan and C. T. Mackown (2002). Copper, manganese and zinc fertilization effects on growth of soybean on a calcareous soil. *Journal of Plant Nutrition* 25:1727-1740.
- Heitholt, J. J., J. J. Sloan, C. T. Mackown and R. I. Cabrera (2003). Soybean growth on calcareous soil as affected by three iron sources. *Journal of Plant Nutrition* 26:935-948.
- Hebborn, C. A., P. Pedas, J. K. Schjoerring, L. Knudsen and S. Husted (2005). Genotypic differences in manganese efficiency: field experiments with winter barley (*Hordeum vulgare* L.). *Plant Soil* 272:233-244.
- Heitholt, J. J., J. J. Sloan and C. T. MacKown (2002). Copper, manganese, and zinc fertilization effects on growth of soybean on a calcareous soil. *J. Plant Nutr.* 25:1727-1740.
- Halliwell, B. and C. Foyer (1978). Properties and physiological function of a glutathione reductase purified from spinach leaves by affinity chromatography. *Planta.* 139:9-17.
- Habibi, D., M. M. Boojar, A. Mahmoudi, M. R. Ardakani and D. Taleghani (2004). Antioxidative enzymes in sunflower subjected to drought stress. 4th International Crop Science Congress, Australia.
- Jiang, Y. and B. Huang (2001). Drought and heat stress injury to two cool-season turf grasses in relation to antioxidant metabolism and lipid peroxidation. *Crop Sci.* 41:436-442.
- Kalbasi, M., F. Filsoof and Y. Rezai-Nejad. (1988). Effect of sulfur treatment on yield and uptake of Fe, Zn and Mn by corn, sorghum and soybean. *J. Plant Nutr.* 11:1353-1360.
- Kaya, C. and D. Higgs (2002). Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. *Sci. Hortic.* 93:53-64.
- Khan, H. R., G. K. McDonald and Z. Rengel (2003). Zn fertilization improves water use efficiency, grain yield and seed Zn content in chickpea. *Plant Soil* , 249:389-400.
- Kheir, N. F., E. Z. Harb, H. A. Moursi and S. H. El-Gayar (1991). Effect of Salinity and Fertilization on Flax Plants (*Linum usitatissimum* L.). II. Chemical Composition, *Bull. Fac. Agric.* 42:57-70.
- Lewis, D. C. and J. D. McFarlane (1986). Effect of foliar applied manganese on the growth of safflower (*Carthamus tinctorious* L.) and the diagnosis of manganese deficiency by plant tissue and seed analysis. *Aus. J. Agric. Res.* 37:567-572.
- Lu, S., X. Liu, L. Li, F. Zhang, X. Zeng and C. Tang (2004). Effect of manganese spatial distribution in the soil profile on wheat growth in rice-wheat rotation. *Plant Soil* 261:39-46.
- Li, D., W. Cong, Z. Cai, D. Shi and F. Ouyang (2002). Response of growth and photosynthesis of marine red tide alga *Heterosigma akashiwo* to iron and iron stress condition. *Biotechnol. Lett.* 24:743-747.
- Marschner, H. (1995). *Mineral Nutrition of Higher Plants.* Academic Press, New York.
- Masoni, A., A. Evacoli and M. Mavoti (1996). Spectral of leaves deficient in iron, sulphur, magnesium and manganese. *Agron. J.* 88(6):937-943.
- Metcalfe, L. C., A. A. Schmitz and J. R. Pelka (1966). Rapid preparation of methyl esters from lipid for gas chromatography analysis. *Analytical chemistry* 38:514-515.
- Mirzapour, M. H. and A. H. Khoshgoftar (2006). Zinc application effects on yield and seed oil content of sunflower

- grown on a saline calcareous soil. J. Plant Nutr. 29:1719-1727.
- Movchan, V. G and I. G. Sobonikova (1972). Zinc fertilizers and the Uptake of Mineral Nutrients by Vinges Growing on the Southern Chozem." Agrokhimiya, 43(5):123-130.
- Nikolic, M. and R. Kastori (2000). Effect of bicarbonate and Fe supply on Fe nutrition of grapevine. J. Plant Nutr. 23:1619-1627.
- Pandey, N., G. C. Pathak and C. P Sharma (2006). Zinc is critically required for pollen function and fertilisation in lentil. J. Trace Elem. Med. Biol. 20: 89-96.
- Sahrawat, K. L. (2000). Elemental composition of the rice plant as affected by iron toxicity under field conditions. Comm. Soil Sci. Plant Anal. 31:2819-2827.
- Sarkar, D., B. Mandal and M. C. Kundu (2007). Increasing use efficiency of boron fertilisers by rescheduling the time and methods of application for crops in India. Plant Soil 301: 77-85.
- Singh, B. P., R. P. Singh and T. P. Yadav (1975). Effect of soil application of zinc yield. Its attributes and oil content of toria (*B. Campestris* var toria). Haryana Agric. University, J. India. 12:283-285.
- Singh, S. (2000). Effect of Fe, Zn on Growth of canola. S. Environmental. 34: 1-2, 57-63.
- Singh, S. and S. Sinha (2005). Accumulation of metals and its effects in *Brassica juncea* (L.) Czern. (cv. Rohini) grown on various amendments of tannery waste, Ecotoxicol. Environ. Safe. 62 (2005), p. 118-127.
- Sultana, N., T. Ikeda and M. A. Kashem (2001). Effect of foliar spray of nutrient solutions on photosynthesis, dry matter accumulation and yield in seawater-stressed rice. Environ. Exp. Bot. 46:129-140.
- Thomas, P., J. B. Mythili and K. S. Shivashankara (2000). Effects of photo oxidative loss of FeNa₂ EDTA and of higher iron supply on chlorophyll content, growth and propagation rate in triploid water melon cultures. In Vitro Cell. Dev. Biol. Plant 36:537-542.
- Welch, R. M., W. H. Allaway, W. A. House and I. Kubota (1991). Geographic distribution of trace element problems. In: Mordvedt, J. J., Cox, F.R.,
- Wissuwa, M., A. M. Ismail and R. D. Graham (2008). Rice grain zinc concentrations as affected by genotype native soil-zinc availability, and zinc fertilization. Plant Soil. 306:37-48.
- Zlatimira, S. and S. Doncheva (2002). The effect of Zn supply and succinate treatment on plant growth and mineral uptake in pea plant. Braz. J. Plant Physiol. 14:111-116.
- Zakaria, M. Sawan, A. Saeb Hafez and A. E. Basyony (2001). Effect of Nitrogen and Zinc Fertilization and Plant Growth Retardants on Cottonseed, Protein, Oil Yields, and Oil Properties. JAOCS, 78(11):18-28.